PubMed		
Display Settings:	Abstract	nature

Nature. 2014 Sep 17. doi: 10.1038/nature13793. [Epub ahead of print]

Artificial sweeteners induce glucose intolerance by altering the gut microbiota.

Suez J¹, Korem T², Zeevi D², Zilberman-Schapira G³, Thaiss CA¹, Maza O¹, Israeli D⁴, Zmora N⁵, Gilad S⁶, Weinberger A⁷, Kuperman Y⁸, Harmelin A⁸, Kolodkin-Gal I⁹, Shapiro H¹, Halpern Z¹⁰, Segal E⁷, Elinav E¹.

Author information

Abstract

Non-caloric artificial sweeteners (NAS) are among the most widely used food additives worldwide, regularly consumed by lean and obese individuals alike. NAS consumption is considered safe and beneficial owing to their low caloric content, yet supporting scientific data remain sparse and controversial. Here we demonstrate that consumption of commonly used NAS formulations drives the development of glucose intolerance through induction of compositional and functional alterations to the intestinal microbiota. These NAS-mediated deleterious metabolic effects are abrogated by antibiotic treatment, and are fully transferrable to germ-free mice upon faecal transplantation of microbiota configurations from NAS-consuming mice, or of microbiota anaerobically incubated in the presence of NAS. We identify NAS-altered microbial metabolic pathways that are linked to host susceptibility to metabolic disease, and demonstrate similar NAS-induced dysbiosis and glucose intolerance in healthy human subjects. Collectively, our results link NAS consumption, dysbiosis and metabolic abnormalities, thereby calling for a reassessment of massive NAS usage.

PMID: 25231862 [PubMed - as supplied by publisher]







LinkOut - more resources

PubMed Commons

PubMed Commons home

0 comments

How to join PubMed Commons

1 of 1 9/25/2014 12:42 PM